

FOR INFORMATION AND ACTION
DA-2018-14
April 25, 2018

Subject: APHIS Adds Seed and Underground Crops to the Exempted Host List for the Light Brown Apple Moth (*Epiphyas postvittana*) in California

To: State and Territory Agricultural Regulatory Officials

Effective immediately, the Animal and Plant Health Inspection Service (APHIS) is adding commercially produced seed with no additional plant material; underground crops with no remaining above ground material; and fruits of litchi tomato, eggplant, and zucchini to the light brown apple moth (LBAM) exempted host list. The exempted host list is included as an attachment to this letter and posted at <http://www.aphis.usda.gov/planthealth/lbam>.


“Exempt host commodities” are exempt from the conditions required in the LBAM Federal Domestic Quarantine Order for interstate movement of regulated articles. The exemption is applicable only to commercially-produced commodities based on the pest mitigations provided through industry standards of production, harvesting, and packaging practices for each of the exempted commodities. Commodities that are not produced using these industry practices remain subject to the program requirements for interstate movement. New commodities added to the list are in **bold** font.

APHIS, in cooperation with the California Department of Food and Agriculture and local county agricultural commissioners, maintains its survey and regulatory framework to prevent the artificial spread of LBAM to other states or trading partners.

LBAM is native to Australia and has established populations in New Zealand, Ireland, the United Kingdom, and Hawaii. The host range for LBAM is broad with more than 250 plant species. A map of the federal quarantine area is at:

http://www.aphis.usda.gov/planthealth/lbam_maps

For further information about the LBAM program, you may contact National Policy Manager Richard Johnson at 301-851-2109.



Osama El-Lissy
Deputy Administrator
Plant Protection and Quarantine

Attachment: LBAM exempted host list

Appendix 1

Light Brown Apple Moth (LBAM) **Host/Commodity List Exempted from Federal Quarantine Order** Alphabetical listing by scientific name

The host commodities, listed below, are exempt from the conditions required in the LBAM Federal Domestic Quarantine Order for interstate movement of regulated articles. The exemption is applicable only to commercially-produced commodities based on the pest mitigations provided through industry standards of production, harvesting, and packaging practices for each of the exempted commodities. Commodities that are not produced using these industry practices remain subject to the program requirements for interstate movement. New commodities added to the list are in **bold font**. **Part A** is agricultural exemptions (pages 1 - 9). **Part B** is horticultural exemptions (nursery stock, cut flower, floral greenery and propagative materials (pages 9 - 12)).

Part A

Host (Scientific Name)	Host (Common Name)	Justification for Exemption
<i>Actinidia spp.</i>	Kiwifruit	Pest management guidelines in California for kiwifruit include visual inspection at prebloom and bloom; chemical application if applicable. Kiwifruit is hand harvested; leaves removed; thoroughly inspected prior to packing.
<i>Allium cepa</i>	Green Onion (with tops cut)	IPM practices implemented by producers, including the use of routine chemical applications that are designed to suppress Lepidoptera pests; also target LBAM. In addition, green onions are hand-picked; tops cut by hand, inspected, washed, and packed.
<i>Allium cepa</i>	Onion, dry	This product is harvested; tops removed; cured in the field prior to storage. Before packing, the onion is thoroughly inspected.

Host (Scientific Name)	Host (Common Name)	Justification for Exemption
<i>Allium sativum</i>	Garlic	This product is harvested; tops removed; cured in the field prior to storage. Before packing, the garlic is thoroughly inspected.
<i>Apium</i> spp.	Celery	IPM practices implemented by producers, including the use of routine chemical applications that are designed to suppress Lepidoptera pests, also target LBAM. Harvest practices focus heavily on visual inspection of leaves in the field before packing and any blemished product does not leave the production site. Leaves are washed and inspected again before packing.
<i>Asparagus officinalis</i>	Asparagus	Only the asparagus spears are harvested. After harvest they are washed, inspected and bundled before packing and shipping. Any blemished spears would be culled in the field.
<i>Beta</i> spp.	Beet	This root crop is generally harvested mechanically with leaves removed in the field. Beets are processed in packing plants where they are washed, brushed, inspected and packaged before shipping.

Host (Scientific Name)	Host (Common Name)	Justification for Exemption
<i>Brassica</i> spp.	Broccoli Brussel sprouts Cauliflower Cabbage Kale Bok Choy Kohlrabi	Integrated Pest Management (IPM) practices implemented by producers, including the use of routine chemical applications that are designed to suppress Lepidoptera pests, also target LBAM. In addition, harvesting and packaging practices provide the safeguard necessary to eliminate the risk of harboring LBAM. As each of the listed <i>Brassica</i> spp. commodities is handpicked, the outer leaves are removed, and inspected for quality, as well as ensuring freedom from pests and diseases before packing and storage. Blemished commodities do not leave the production area and are either disked into the ground or consumed locally.
<i>Brassica</i> spp.	Mustard Collards	In addition to the above practices, mustard leaves and collards are carefully washed and inspected again prior to packing and storage.
<i>Capsicum</i> spp.	Peppers	Only the peppers are harvested (no leaves) and the product is inspected thoroughly before shipping from the field.
<i>Citrullus lanatus</i> <i>Citrullus vulgaris</i>	Watermelon	This product is harvested without leaves; visually inspected prior to packing.
<i>Citrus</i> spp.	Citrus fruit (conventionally produced)	LBAM survival on citrus is low compared with non-citrus hosts. The combination of low field prevalence and packing procedures make it highly unlikely that a foundation LBAM population could be moved out of the quarantine area with citrus fruit after conventional production and harvesting.
<i>Coriander sativum</i>	Cilantro	IPM practices implemented by producers, including the routine use of chemical applications that are designed to suppress Lepidoptera pests; also target LBAM; product is handpicked; inspected; hydro-cooled.
<i>Cucumis melo</i>	Melons	IPM practices implemented by producers, including routine use of chemical applications that are designed to suppress Lepidoptera pests; also target LBAM; product is handpicked; inspected prior to packing.

Host (Scientific Name)	Host (Common Name)	Justification for Exemption
<i>Cucumis sativus</i>	Cucumbers	LBAM does not appear to be a pest of cucumbers. Additionally, IPM practices implemented by producers would minimize risk of LBAM survival; fruit is hand harvested; visually inspected prior to packing; stored and shipped at low temperatures.
<i>Cucurbita</i> spp.	Squash Pumpkin	Only the squash and pumpkin are harvested (no leaves) and the product is inspected thoroughly before shipping from the field.
<i>Cynara scolymus</i>	Globe artichoke	Only artichoke thistle flower heads are harvested and are inspected twice before packing and shipping from the field to the cooler. They are inspected again at the cooler before final packing and shipping to market.
<i>Daucus carota</i>	Carrot	IPM practices implemented by producers, including routine use of chemical applications that are designed to suppress Lepidoptera pests; also target LBAM. Product is handpicked, inspected and washed prior to packing.
<i>Diospyros kaki</i>	Persimmon	Pest management in California for persimmon, including hand harvesting, leave removal, and visual inspection, minimize the likelihood of LBAM movement.
<i>Foeniculum vulgare</i>	Fennel	IPM practices implemented by producers, including routine use of chemical applications that are designed to suppress Lepidoptera pests; also target LBAM. Product is handpicked; inspected prior to packing.
<i>Fragaria x ananassa</i>	Strawberry (conventionally produced)	Based on evidence and our current understanding of LBAM biology, we have determined that it is appropriate to remove picked-and-packed fresh strawberry fruit produced using conventional (non-organic) field-production methodologies.
<i>Fragaria</i> spp.	Strawberry (dormant, frozen strawberry nursery stock)	The product is dormant, frozen strawberry nursery stock grown by State certified producers.

Host (Scientific Name)	Host (Common Name)	Justification for Exemption
<i>Gossypium hirsutum</i> <i>Gossypium barbadense</i>	Cotton	IPM practices implemented by producers include routine use of chemical applications that are effective on LBAM. Product is harvested mechanically; heated; ginned; and compressed into bales.
<i>Juglans</i> spp.	Walnuts	All commercial walnuts are mechanically harvested free of husks and leaves and are taken in bins to packing plants where they are inspected thoroughly before packing.
<i>Lactuca sativa</i>	Head Lettuce	IPM practices implemented by producers, including the use of routine chemical applications that are designed to suppress Lepidoptera pests, also target LBAM. In addition, harvesting and packaging practices provide the safeguard necessary to eliminate the risk of harboring LBAM. Each head of lettuce is handpicked; the outer leaves are removed, and inspected for quality, as well as ensuring freedom from pests and diseases before packing and storage. Blemished lettuce heads do not leave the production site and are either disked into the ground or consumed locally.
<i>Lactuca sativa</i>	Leaf Lettuce	IPM practices implemented by producers, including the use of routine chemical applications that are designed to suppress Lepidoptera pests, also target LBAM. In addition, leaf lettuce is mechanically harvested, thoroughly washed, and then inspected before shipping from the regulated area.
<i>Malus domestica</i>	Apple	Apple producer best management practices greatly reduce both the prevalence of LBAM in production areas and the likelihood that LBAM will be present on market-ready commodities. Movement of commercially produced apples represents a low risk pathway for establishment of new LBAM infestations after conventional production and harvesting.
<i>Olea</i> spp.	Olive	The product is handpicked and placed in bins that are immediately taken to packing plants where the olives are placed in a brine solution for curing.

Host (Scientific Name)	Host (Common Name)	Justification for Exemption
<i>Opuntia</i> spp.	Tuna (fruit and pad) (Prickly pear)	This product is handpicked, allowing for in-field inspection. In addition to this, processing and packaging practices involving the use of hot water dip, spine removal, and post-harvest storage provides the safeguard necessary to eliminate the risk of harboring or reinfestation of LBAM.
<i>Persea americana</i>	Avocado (conventionally and organically grown)	Analysis of commercial production practices determined avocados present a low risk of dispersing viable populations of the pest. Standard practices include thorough inspection and hand packing of each harvested fruit, with damaged fruit culled. The commercial processing would result in the detection and removal of any leaves attached to the fruit, any larvae or pupae hidden underneath the leaves, and any larvae or pupae present without leaves.
<i>Petroselinum crispum</i>	Parsley	IPM practices implemented by producers, including the use of routine chemical applications that are designed to suppress Lepidoptera pests, also target LBAM. Harvest practices focus heavily on visual inspection of leaves in the field before packing and any blemished product does not leave the production site. Leaves are washed and inspected again before packing.
<i>Phaseolus vulgaris</i>	Green beans	IPM practices implemented by producers including chemical application greatly reduce LBAM if present in production areas. Beans are either machine or hand harvested; visually inspected prior to packing.
<i>Pistacia vera</i>	Pistachio	Commercial pistachios are mechanically harvested; dried; leaves removed; visual inspection during packing.
<i>Prunus amygdalus</i> <i>Prunus dulcis</i>	Almond	Commercial almonds are mechanically harvested; dried; fumigated. Leaves and hulls are removed; visual inspection of nuts prior to packing.

Host (Scientific Name)	Host (Common Name)	Justification for Exemption
<i>Prunus avium</i>	Cherry	The combination of low field prevalence and packing procedures make it highly unlikely that a foundation LBAM population could be moved out of the quarantine area with cherry fruit after conventional production and harvesting.
<i>Prunus persica</i> , <i>P. armeniaca</i> , <i>P. persica</i> var. <i>nucipersica</i> , <i>P. persica</i> , <i>P. domestica</i> or <i>P. salicina</i> , <i>P. armeniaca</i> x <i>P.</i> <i>domestica</i> x <i>P.</i> <i>armeniaca</i> , <i>P. armeniaca</i> x <i>P.</i> <i>domestica</i> x <i>P.</i> <i>domestica</i>	Stone fruit (peaches, plums, nectarines, and apricots)	IPM practices implemented by producers, including routine chemical applications to suppress Lepidoptera pests, keep LBAM prevalence very low in stone fruit production areas. The combination of low field prevalence and packing procedures (i.e. hand packing and refrigeration at 1°C prior to shipment) make it highly unlikely that a foundation LBAM population could be moved out of the quarantine area with stone fruit.
<i>Punica granatum</i>	Pomegranate	This product is harvested without leaves; washed; brushed; waxed; inspected during packing.
<i>Pyrus communis</i>	Pear	Attributes of pears and the industry practices, such as monitoring for pests, application of mating disruption, visual inspection, and hand-packing without leaves provide further mitigations to prevent human assisted dispersal of LBAM.
<i>Rhaphanus sativus</i>	Daikon	IPM practices implemented by producers including chemical application greatly reduce LBAM if present in production areas. Daikon, with or without tops, is hand-picked; washed; and visually inspected prior to packing.
<i>Raphanus</i> spp.	Radish (w/o tops)	This root crop is generally harvested mechanically with leaves removed in the field. Radishes are processed through packing plants where they are washed, brushed, inspected and packaged before shipping.

Host (Scientific Name)	Host (Common Name)	Justification for Exemption
<i>Spinacia oleracea</i>	Spinach	IPM practices implemented by producers, including the use of routine chemical applications that are designed to suppress Lepidoptera pests, also target LBAM. In addition, spinach is mechanically harvested, thoroughly washed, and then inspected before shipping from the regulated area.
<i>Solanum lycopersicum</i>	Tomato	Only the tomatoes are harvested (no leaves) and the product is inspected thoroughly before shipping from the field.
<i>Solanum melongena</i>	Eggplant	IPM practices implemented by producers would minimize risk of LBAM survival; fruit is hand harvested; visually inspected prior to packing; stored and shipped at low temperatures.
<i>Solanum</i> spp.	Potato	This root crop is generally harvested mechanically with no leaves attached. Potatoes are processed in packing plants where they are washed, brushed, inspected and packaged before shipping.
<i>Taraxacum officinale</i>	Dandelion green	IPM practices implemented by producers, including routine use of chemical applications that are designed to suppress Lepidoptera pests; also target LBAM. Product is handpicked; inspected prior to packing.
<i>Vaccinium</i> spp.	Blueberry	IPM practices implemented by producers including monitoring and surveillance greatly reduce LBAM if present; fruit is hand harvested; leaves removed; visually inspected prior to packing.
Various species of forage, no scientific name provided	Baled Hay	This product is generally harvested mechanically, dried, and compressed into bales.
Various species of fruit, no scientific name provided	Dried Fruit (such as raisins, dates, etc)	Analysis of scientific literature showed that LBAM is not a storage pest.
<i>Vitis vinifera</i>	Table grapes	IPM practices implemented by producers minimize risk of LBAM; fruit is hand harvested; visually inspected prior to packing; stored and shipped at low temperatures.

Host (Scientific Name)	Host (Common Name)	Justification for Exemption
<i>Vitis vinifera</i>	Wine grapes	IPM practices implemented by producers minimize risk of LBAM; fruit is harvested in a manner that is not conducive to maintaining live larvae, visually inspected prior to packing, stored and shipped at low temperatures. Green waste, the product left after the processing of wine grapes, is simply the next step in the same pathway, thus green waste is also exempt.
<i>Zea mays</i>	Sweet Corn	IPM practices implemented by producers, including the use of routine chemical applications that are designed to suppress Lepidoptera pests, also target LBAM. In addition to this, processing and packaging practices involving the use of an ice water bath immediately followed by the commodity being packed in ice for shipment provides the safeguard necessary to eliminate the risk of harboring or reinfestation of LBAM.
<i>Ziziphus jujube</i>	Jujube	Fruit is hand harvested without leaves; inspected prior to packing.
<i>Brassica rapa rapa</i>	Turnips (w/o tops)	Analysis of scientific literature showed that LBAM are not associated with harvested commodity.
<i>Pastinaca sativa</i>	Parsnips (w/o tops)	Analysis of scientific literature showed that LBAM are not associated with harvested commodity.
<i>Brassica napobrassica</i>	Rutabaga	Analysis of scientific literature showed that LBAM are not associated with harvested commodity.
Various	Seed crops without any plant material	Analysis of scientific literature showed that LBAM are not associated with harvested commodity.
<i>Solanum sisymbriifolium</i>	Litchi tomato (harvested as a horticultural fruit)	IPM practices implemented by producers, including the use of routine chemical applications that are designed to suppress Lepidoptera pests, also target LBAM. In addition to this, processing and packaging practices provides the safeguard necessary to eliminate the risk of harboring LBAM.

<i>Solanum melongena</i>	Eggplant (harvested as a horticultural fruit)	IPM practices implemented by producers, including the use of routine chemical applications that are designed to suppress Lepidoptera pests, also target LBAM. In addition to this, processing and packaging practices provides the safeguard necessary to eliminate the risk of harboring LBAM.
<i>Cucurbita pepo</i> var. <i>cylindrical</i>	Zucchini (harvested as a horticultural fruit)	IPM practices implemented by producers, including the use of routine chemical applications that are designed to suppress Lepidoptera pests, also target LBAM. In addition to this, processing and packaging practices provides the safeguard necessary to eliminate the risk of harboring LBAM.

Part B**Horticultural Exemptions: Nursery stock, cut flowers, floral greenery and propagative materials**

Host (Scientific Name)	Host (Common Name)	Justification for Exemption
Cactaceae	Commercially grown cacti	Eligible due to non-host status.
Cycadales	Commercially grown cycads	Eligible due to non-host status.
<i>Fragaria</i> spp.	Strawberry (dormant, frozen strawberry nursery stock)	The product is dormant, frozen strawberry nursery stock grown by State certified producers.
Palmaceae	Commercially grown palms	Eligible due to non-host status.
Succulents: Agavaceae - <i>Mangave</i> Aizoaceae- <i>Aloinopsis</i> , <i>Antimima</i> , <i>Aptenia</i> , <i>Argyroderma</i> , <i>Carpobrotus</i> , <i>Cephalophyllum</i> , <i>Cerochlamys</i> , <i>Conophytum</i> , <i>Cylindrophyllum</i> ,	Various genera of commercially grown succulents, as provided. With the exception of succulents of the genera <i>Senecio</i> and <i>Euphorbia</i> .	Eligible due to non-host status. Possible host associations with members of the genera <i>Senecio</i> and <i>Euphorbia</i> do not allow those plants to be exempted at this time. These commodities pose no or very low risk of spreading LBAM from infested areas due to the morphological nature of the plants, the general and LBAM specific integrated pest management (IPM) practiced in nurseries and the nature of the commercial trade of cacti and succulents.

<p><i>Delosperma,</i> <i>Dinteranthus,</i> <i>Drosantemum,</i> <i>Faucaria,</i> <i>Fenestriaria, Frithia,</i> <i>Gibbaem,</i> <i>Glottiphyllum,</i> <i>Lampranthus,</i> <i>Lapidaria, Lithops,</i> <i>Oscularia, Pleiospilos,</i> <i>Rabiea,</i> <i>Rhombophyllum,</i> <i>Ruschia, Schwantesia,</i> <i>Titanopsis, and</i> <i>Trichodiadema.</i></p> <p>Aloaceae – <i>Gasteraloe,</i> <i>Gasterworthia</i></p> <p>Amaryllidaceae - <i>Allium</i></p> <p>Apocynaceae - <i>Caralluma,</i> <i>Ceropegia,</i> <i>Cynanchum, Fockea,</i> <i>Hoodia, Huernia,</i> <i>Orbea, Stapelia</i></p> <p>Asparagaceae - <i>Agave,</i> <i>Aloe, Bowiea,</i> <i>Calibanus, Dasyilirion,</i> <i>Drimiopsis, Furcraea,</i> <i>Gasteria, Haworthia,</i> <i>Hesperaloe,</i> <i>Ledebouria,</i> <i>Ornithogalum,</i> <i>Sansevierea,</i> <i>Veltheimia, Yucca</i></p> <p>Commelinaceae - <i>Tradescantia</i></p>		
--	--	--

<p>Compositae - <i>Kleinia</i>, <i>Othonna</i></p> <p>Crassulaceae - <i>Adromischus</i>, <i>Aeonium</i>, <i>Aichryson</i>, <i>Bryophyllum</i>, <i>Cotyledon</i>, <i>Cotyledon</i>, <i>Crassula</i>, <i>Crassula</i>, <i>Cremnosedum</i>, <i>Dudleya</i>, <i>Echeveria</i>, <i>Graptopetalum</i>, <i>Graptosedum</i>, <i>Graptoveria</i>, <i>Jovibarba</i>, <i>Kalanchoe</i>, <i>Monanthes</i>, <i>Orostachys</i>, <i>Pachyphytum</i>, <i>Pachysedum</i>, <i>Pachyveria</i>, <i>Rosularia</i>, <i>Sedeveria</i>, <i>Sedum</i>, <i>Sempervivella</i>, <i>Sempervivum</i>, <i>Tacitus</i></p> <p>Cucurbitaceae - <i>Xerosycos</i></p> <p>Didiereaceae - <i>Alluaudia</i>, <i>Didierea</i></p> <p>Dioscoreaceae - <i>Dioscorea</i>, <i>Testudinaria</i></p> <p>Euphorbiaceae - <i>Monadenium</i>, <i>Pedilanthus</i>, <i>Synadenium</i>.</p> <p>Fouquieriaceae - <i>Fouquieria</i></p> <p>Gesneriaceae - <i>Reichsteinaria</i></p>		
--	--	--

<p>Malvaceae – <i>Bombax</i></p> <p>Piperaceae - <i>Peperomia</i></p> <p>Portulacaceae - <i>Anacampseros</i>, <i>Calandrinia</i>, <i>Portulaca</i>, <i>Portulacaria</i></p> <p>Vitaceae - <i>Cyphostemma</i></p> <p>Xanthorrhoeaceae - <i>Bulbine</i></p>		
<p>Host (Scientific Name)</p>	<p>Host (Common Name)</p>	<p>Justification for Exemption</p>
<p>Various</p>	<p>Propagative material: Bulbs, tissue cultivated plantlets callused vegetative cuttings, seedlings, divisions, unrooted vegetative cuttings, callused vegetative cuttings, rooted vegetative cuttings, under condition that the plants do not exceed 10 cm in height.</p>	<p>These commodities pose no or very low risk of spreading LBAM from infested areas due to the morphological nature of the plants, the general and LBAM specific integrated pest management (IPM) practices in nurseries and the nature of the commercial trade of propagative material. Further definitions of the plant material are provided in the Pest Risk Evaluation of propagative material.</p>
<p>Various</p>	<p>Cut flowers and floral greenery (commercially produced)</p>	<p>IPM practices and industry standards for handling commercially produced cut flowers and floral greenery, minimize the likelihood of LBAM movement.</p>