

FOR INFORMATION AND ACTION
DA-2009-41
August 21, 2009

SUBJECT: Mediterranean Fruit Fly, *Ceratitis capitata* – Establishment of a Third Quarantine Area in San Diego County, California

TO: STATE AND TERRITORY AGRICULTURAL REGULATORY OFFICIALS

The Animal and Plant Health Inspection Service (APHIS) has established a new Mediterranean fruit fly (Medfly) quarantine area in the Imperial Beach area of San Diego County, California, following the confirmed detection of a Medfly population in that area. APHIS is applying restrictions on the interstate movement of regulated articles from the quarantine area. These actions are necessary to prevent the spread of Medfly to noninfested areas of the United States. This Medfly outbreak is considered to be transient, actionable, and under eradication.

On July 30, 2009, one adult mated female Medfly was detected on a residential property in the Imperial Beach area of San Diego, California. APHIS, in cooperation with the California Department of Food and Agriculture and the San Diego County Department of Agriculture, is responding to this confirmed Medfly finding. This confirmed finding triggered the establishment of the new quarantine area, which encompasses approximately 37 square miles of San Diego County. This new, distinct Medfly quarantine area is southwest of two previously established Medfly quarantine areas in San Diego County. The southern border of this newest Medfly quarantine area ends at the international border with Mexico. There is no commercial production of host commodities within this new quarantine area.

Fruit fly traps have been deployed at protocol levels to conduct a delimitation survey surrounding the new Medfly detection site. APHIS; in cooperation with officials from Mexico's Department of Agriculture, Livestock, Rural Development, Fisheries and Food; are also increasing trapping activities in Mexico along the international border with the United States. Spinosad foliar bait spray treatments are being applied to all host trees within 200 meters of the detection site at 7 to 10 day intervals. As a treatment to control Medfly populations within the new quarantine area, Sterile Insect Technique (SIT) will be conducted in a 7.19 square mile treatment area surrounding the detection site. SIT is an eradication tool, which consists of the aerial release of sterile male Medflies to inundate the detected wild population to reduce reproduction. Sterile male Medflies are released at rate of 250,000 per square mile per week. The weekly release of Sterile Medflies will continue in the San Diego area through two projected Medfly life cycles since the date of the latest detection.

This action became effective on August 13, 2009. This establishment of this quarantine area will be reflected on the following designated website, which contains a description of all the current Federal fruit fly quarantine areas:

http://www.aphis.usda.gov/plant_health/plant_pest_info/fruit_flies/index.shtml

APHIS anticipates following this action with the publication of a notice in the Federal Register informing the public of this change. For additional information on this Medfly Eradication program, you may contact Wayne Burnett, APHIS Exotic Fruit Fly Director, at (301) 734-6553.

/s/ John H. Payne for

Rebecca A. Bech
Deputy Administrator
Plant Protection and Quarantine